

Sixth Term Examination Papers

9470

MATHEMATICS 2

Morning

WEDNESDAY 19 JUNE 2013

Time: 3 hours

* 2 1 2 7 1 6 4 7 7 1 *

Additional Materials: Answer Booklet
Formulae Booklet

INSTRUCTIONS TO CANDIDATES

Please read this page carefully, but do not open this question paper until you are told that you may do so.

Write your name, centre number and candidate number in the spaces on the answer booklet.

Begin each answer on a new page.

Write the numbers of the questions you answer in the order attempted on the front of the answer booklet.

INFORMATION FOR CANDIDATES

Each question is marked out of 20. There is no restriction of choice.

All questions attempted will be marked.

Your final mark will be based on the **six** questions for which you gain the highest marks.

You are advised to concentrate on no more than **six** questions. Little credit will be given for fragmentary answers.

You are provided with a Mathematical Formulae Booklet.

Calculators are not permitted.

Please wait to be told you may begin before turning this page.

This question paper consists of 8 printed pages and 4 blank pages.

Section A: Pure Mathematics

- 1** (i) Find the value of m for which the line $y = mx$ touches the curve $y = \ln x$.
If instead the line intersects the curve when $x = a$ and $x = b$, where $a < b$, show that $a^b = b^a$. Show by means of a sketch that $a < e < b$.
- (ii) The line $y = mx + c$, where $c > 0$, intersects the curve $y = \ln x$ when $x = p$ and $x = q$, where $p < q$. Show by means of a sketch, or otherwise, that $p^q > q^p$.
- (iii) Show by means of a sketch that the straight line through the points $(p, \ln p)$ and $(q, \ln q)$, where $e \leq p < q$, intersects the y -axis at a positive value of y . Which is greater, π^e or e^π ?
- (iv) Show, using a sketch or otherwise, that if $0 < p < q$ and $\frac{\ln q - \ln p}{q - p} = e^{-1}$, then $q^p > p^q$.

- 2** For $n \geq 0$, let

$$I_n = \int_0^1 x^n(1-x)^n dx.$$

- (i) For $n \geq 1$, show by means of a substitution that

$$\int_0^1 x^{n-1}(1-x)^n dx = \int_0^1 x^n(1-x)^{n-1} dx$$

and deduce that

$$2 \int_0^1 x^{n-1}(1-x)^n dx = I_{n-1}.$$

Show also, for $n \geq 1$, that

$$I_n = \frac{n}{n+1} \int_0^1 x^{n-1}(1-x)^{n+1} dx$$

and hence that $I_n = \frac{n}{2(2n+1)} I_{n-1}$.

- (ii) When n is a positive integer, show that

$$I_n = \frac{(n!)^2}{(2n+1)!}.$$

- (iii) Use the substitution $x = \sin^2 \theta$ to show that $I_{\frac{1}{2}} = \frac{\pi}{8}$, and evaluate $I_{\frac{3}{2}}$.

3 (i) Given that the cubic equation $x^3 + 3ax^2 + 3bx + c = 0$ has three distinct real roots and $c < 0$, show with the help of sketches that either exactly one of the roots is positive or all three of the roots are positive.

(ii) Given that the equation $x^3 + 3ax^2 + 3bx + c = 0$ has three distinct real positive roots show that

$$a^2 > b > 0, \quad a < 0, \quad c < 0. \quad (*)$$

[Hint: Consider the turning points.]

(iii) Given that the equation $x^3 + 3ax^2 + 3bx + c = 0$ has three distinct real roots and that

$$ab < 0, \quad c > 0,$$

determine, with the help of sketches, the signs of the roots.

(iv) Show by means of an explicit example (giving values for a , b and c) that it is possible for the conditions (*) to be satisfied even though the corresponding cubic equation has only one real root.

4 The line passing through the point $(a, 0)$ with gradient b intersects the circle of unit radius centred at the origin at P and Q , and M is the midpoint of the chord PQ . Find the coordinates of M in terms of a and b .

(i) Suppose b is fixed and positive. As a varies, M traces out a curve (the *locus* of M). Show that $x = -by$ on this curve. Given that a varies with $-1 \leq a \leq 1$, show that the locus is a line segment of length $2b/(1 + b^2)^{\frac{1}{2}}$. Give a sketch showing the locus and the unit circle.

(ii) Find the locus of M in the following cases, giving in each case its cartesian equation, describing it geometrically and sketching it in relation to the unit circle:

(a) a is fixed with $0 < a < 1$, and b varies with $-\infty < b < \infty$;

(b) $ab = 1$, and b varies with $0 < b \leq 1$.

- 5 (i) A function $f(x)$ satisfies $f(x) = f(1-x)$ for all x . Show, by differentiating with respect to x , that $f'(\frac{1}{2}) = 0$. If, in addition, $f(x) = f(\frac{1}{x})$ for all (non-zero) x , show that $f'(-1) = 0$ and that $f'(2) = 0$.

- (ii) The function f is defined, for $x \neq 0$ and $x \neq 1$, by

$$f(x) = \frac{(x^2 - x + 1)^3}{(x^2 - x)^2}.$$

Show that $f(x) = f(\frac{1}{x})$ and $f(x) = f(1-x)$.

Given that it has exactly three stationary points, sketch the curve $y = f(x)$.

- (iii) Hence, or otherwise, find all the roots of the equation $f(x) = \frac{27}{4}$ and state the ranges of values of x for which $f(x) > \frac{27}{4}$.

Find also all the roots of the equation $f(x) = \frac{343}{36}$ and state the ranges of values of x for which $f(x) > \frac{343}{36}$.

- 6 In this question, the following theorem may be used.

Let u_1, u_2, \dots be a sequence of (real) numbers. If the sequence is bounded above (that is, $u_n \leq b$ for all n , where b is some fixed number) and increasing (that is, $u_n \geq u_{n-1}$ for all n), then the sequence tends to a limit (that is, converges).

The sequence u_1, u_2, \dots is defined by $u_1 = 1$ and

$$u_{n+1} = 1 + \frac{1}{u_n} \quad (n \geq 1). \quad (*)$$

- (i) Show that, for $n \geq 3$,

$$u_{n+2} - u_n = \frac{u_n - u_{n-2}}{(1 + u_n)(1 + u_{n-2})}.$$

- (ii) Prove, by induction or otherwise, that $1 \leq u_n \leq 2$ for all n .

- (iii) Show that the sequence u_1, u_3, u_5, \dots tends to a limit, and that the sequence u_2, u_4, u_6, \dots tends to a limit. Find these limits and deduce that the sequence u_1, u_2, u_3, \dots tends to a limit.

Would this conclusion change if the sequence were defined by (*) and $u_1 = 3$?

- 7 (i) Write down a solution of the equation

$$x^2 - 2y^2 = 1, \quad (*)$$

for which x and y are non-negative integers.

Show that, if $x = p$, $y = q$ is a solution of $(*)$, then so also is $x = 3p + 4q$, $y = 2p + 3q$. Hence find two solutions of $(*)$ for which x is a positive odd integer and y is a positive even integer.

- (ii) Show that, if x is an odd integer and y is an even integer, $(*)$ can be written in the form

$$n^2 = \frac{1}{2}m(m+1),$$

where m and n are integers.

- (iii) The positive integers a , b and c satisfy

$$b^3 = c^4 - a^2,$$

where b is a prime number. Express a and c^2 in terms of b in the two cases that arise.

Find a solution of $a^2 + b^3 = c^4$, where a , b and c are positive integers but b is not prime.

- 8 The function f satisfies $f(x) > 0$ for $x \geq 0$ and is strictly decreasing (which means that $f(b) < f(a)$ for $b > a$).

- (i) For $t \geq 0$, let $A_0(t)$ be the area of the largest rectangle with sides parallel to the coordinate axes that can fit in the region bounded by the curve $y = f(x)$, the y -axis and the line $y = f(t)$. Show that $A_0(t)$ can be written in the form

$$A_0(t) = x_0 (f(x_0) - f(t)),$$

where x_0 satisfies $x_0 f'(x_0) + f(x_0) = f(t)$.

- (ii) The function g is defined, for $t > 0$, by

$$g(t) = \frac{1}{t} \int_0^t f(x) dx.$$

Show that $tg'(t) = f(t) - g(t)$.

Making use of a sketch show that, for $t > 0$,

$$\int_0^t (f(x) - f(t)) dx > A_0(t)$$

and deduce that $-t^2 g'(t) > A_0(t)$.


- (iii) In the case $f(x) = \frac{1}{1+x}$, use the above to establish the inequality

$$\ln \sqrt{1+t} > 1 - \frac{1}{\sqrt{1+t}},$$

for $t > 0$.

Section B: Mechanics

- 9 The diagram shows three identical discs in equilibrium in a vertical plane. Two discs rest, not in contact with each other, on a horizontal surface and the third disc rests on the other two. The angle at the upper vertex of the triangle joining the centres of the discs is 2θ .



The weight of each disc is W . The coefficient of friction between a disc and the horizontal surface is μ and the coefficient of friction between the discs is also μ .

- (i) Show that the normal reaction between the horizontal surface and a disc in contact with the surface is $\frac{3}{2}W$.
 - (ii) Find the normal reaction between two discs in contact and show that the magnitude of the frictional force between two discs in contact is $\frac{W \sin \theta}{2(1 + \cos \theta)}$.
 - (iii) Show that if $\mu < 2 - \sqrt{3}$ there is no value of θ for which equilibrium is possible.
- 10 A particle is projected at an angle of elevation α (where $\alpha > 0$) from a point A on horizontal ground. At a general point in its trajectory the angle of elevation of the particle from A is θ and its direction of motion is at an angle ϕ above the horizontal (with $\phi \geq 0$ for the first half of the trajectory and $\phi \leq 0$ for the second half).

Let B denote the point on the trajectory at which $\theta = \frac{1}{2}\alpha$ and let C denote the point on the trajectory at which $\phi = -\frac{1}{2}\alpha$.

- (i) Show that, at a general point on the trajectory, $2 \tan \theta = \tan \alpha + \tan \phi$.
- (ii) Show that, if B and C are the same point, then $\alpha = 60^\circ$.
- (iii) Given that $\alpha < 60^\circ$, determine whether the particle reaches the point B first or the point C first.

- 11** Three identical particles lie, not touching one another, in a straight line on a smooth horizontal surface. One particle is projected with speed u directly towards the other two which are at rest. The coefficient of restitution in all collisions is e , where $0 < e < 1$.
- (i) Show that, after the second collision, the speeds of the particles are $\frac{1}{2}u(1-e)$, $\frac{1}{4}u(1-e^2)$ and $\frac{1}{4}u(1+e)^2$. Deduce that there will be a third collision whatever the value of e .
- (ii) Show that there will be a fourth collision if and only if e is less than a particular value which you should determine.

Section C: Probability and Statistics

- 12** The random variable U has a Poisson distribution with parameter λ . The random variables X and Y are defined as follows.

$$X = \begin{cases} U & \text{if } U \text{ is } 1, 3, 5, 7, \dots \\ 0 & \text{otherwise} \end{cases}$$
$$Y = \begin{cases} U & \text{if } U \text{ is } 2, 4, 6, 8, \dots \\ 0 & \text{otherwise} \end{cases}$$

- (i) Find $E(X)$ and $E(Y)$ in terms of λ , α and β , where

$$\alpha = 1 + \frac{\lambda^2}{2!} + \frac{\lambda^4}{4!} + \dots \quad \text{and} \quad \beta = \frac{\lambda}{1!} + \frac{\lambda^3}{3!} + \frac{\lambda^5}{5!} + \dots$$

- (ii) Show that

$$\text{Var}(X) = \frac{\lambda\alpha + \lambda^2\beta}{\alpha + \beta} - \frac{\lambda^2\alpha^2}{(\alpha + \beta)^2}$$

and obtain the corresponding expression for $\text{Var}(Y)$. Are there any non-zero values of λ for which $\text{Var}(X) + \text{Var}(Y) = \text{Var}(X + Y)$?

- 13** A biased coin has probability p of showing a head and probability q of showing a tail, where $p \neq 0$, $q \neq 0$ and $p \neq q$. When the coin is tossed repeatedly, runs occur. A *straight run* of length n is a sequence of n consecutive heads or n consecutive tails. An *alternating run* of length n is a sequence of length n alternating between heads and tails. An alternating run can start with either a head or a tail.

Let S be the length of the longest straight run beginning with the first toss and let A be the length of the longest alternating run beginning with the first toss.

- (i) Explain why $P(A = 1) = p^2 + q^2$ and find $P(S = 1)$. Show that $P(S = 1) < P(A = 1)$.
- (ii) Show that $P(S = 2) = P(A = 2)$ and determine the relationship between $P(S = 3)$ and $P(A = 3)$.
- (iii) Show that, for $n > 1$, $P(S = 2n) > P(A = 2n)$ and determine the corresponding relationship between $P(S = 2n + 1)$ and $P(A = 2n + 1)$. [You are advised *not* to use $p + q = 1$ in this part.]

BLANK PAGE

BLANK PAGE

BLANK PAGE

BLANK PAGE